

ANTIPSYCHOTIC MONITORING

	Baseline	At 4-8 Weeks	First 12 Weeks	Every 3 Months	Every 6 Months	Annually	As Clinically Indicated
General Physical Assmt: BP, HR (T, RR as clinically indicated)							
Weight/BMI *							
Lifestyle Assessment: smoking, drug, alcohol, diet, exercise							
Waist Circumference at level of umbilicus							
Fasting Blood Glucose: more often if sig risk factors							
Lipid Screening							
Review of All Medications: including OTC and herbal supp							
Pregnancy Test: test, reproductive status, contraceptive use							
Sexual Function Inquiry: menstrual/libido/erectile disturbance							
EPS including Akathisia							
Tardive dyskinesia: AIMS, more frequent for high risk/elderly							
Prolactin Level							
Ocular Exam: chlorpromazine, prochlorperazine, quetiapine *							
Clozapine CBC Monitoring							
Cardiac Evaluation **							
Assess side effects, symptoms, adherence to treatment plan							

* refer for slit-lamp exam at initiation and six month interval

** assess risk factors, baseline EKG, monitor for symptom of prolonged QT such as syncope

NOTE: THIS REPRESENTS MINIMUM RECOMMENDED MONITORING AND FURTHER EVALUATION AND TESTING MAY BE INDICATED BASED ON THE PROVIDERS CLINICAL JUDGEMENT